



Severe Respiratory Disease Concurrent with the Circulation of H₁N₁ Influenza

Time Series of Confirmed Cases of Influenza in the Federal District and in All of Mexico, According to Month (January 2005-April 2009)




Severe Respiratory Disease Concurrent with the Circulation of H₁N₁ Influenza

Percentage Distribution of Deaths from **Severe Pneumonia** during the 2009 Study Period, as Compared with Influenza Seasons from 2006 through 2008, in Mexico, According to Age Group



Severe Respiratory Disease Concurrent with the Circulation of H₁N₁ Influenza

Percentage Distribution of Illness from **Severe Pneumonia** during the 2009 Study Period, as Compared with Influenza Seasons from 2005 through 2008, in Mexico, According to Age Group



Chowell G et al. N Engl J Med 2009;10.1056/NEJMoa0904023

Severe Respiratory Disease Concurrent with the Circulation of H₁N₁ Influenza

Table 1. Proportion of Mortality from Severe Pneumonia According to Age Group during the 2009 Study Period, as Compared with Influenza Seasons from 2006 through 2008, in Mexico.*

Age Group yr	2006–2008 Influenza Seasons	Mortality March 24 to April 29, 2009	Ratio, Study Period to Referent Period
	percent		
0–4	17	5	0.3
5–9	1	5	7.6
10–14	1	6	10.7
15–19	1	7	8.6
20–24	1	13	11.7
25–29	1	11	10.3
30–34	1	9	6.0
35–39	2	15	9.4
40–44	2	9	4.8
45–49	2	5	2.1
50–54	2	4	1.6
55–59	3	3	1.2
60–64	3	0	0.0
65–69	5	1	0.2
70–74	7	2	0.3
75–79	9	2	0.2
≥80	42	3	0.1

* Data were reported by the National System of Health Care Information and the National Epidemiological Surveillance System (SINAVE).

Proportion of Mortality from Severe Pneumonia According to Age Group during the 2009 Study Period, as Compared with Influenza Seasons from 2006 through 2008, in Mexico

Chowell G et al. N Engl J Med 2009;10.1056/NEJMoa0904023